



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Hence triangle $EC'C' < \text{triangle } EBB'$. But, $ABCD = AB'EC'D + ABB' + EBB'$ and $AB'C'D = AB'ECD + DCC' + EC'C'$.

Hence $ABCD$ is greater than $AB'C'D$.

COROLLARY. It follows, that the quadrilateral of three equal sides, and maximum area, is a trapezoid; that the angles including the fourth side are also equal; that the opposite angles are supplementary; and that the trapezoid is inscriptible.

[To be continued.]

PROFESSOR SYLVESTER'S RECIPROCANTS.

By F. P. MATZ, M. Sc., Ph. D., New Windsor, Maryland.

To those functions of the successive derivatives of y with respect to x , which preserve their form unaltered, except for $dy \propto dx$ as a factor, when the independent and dependent variables x and y are interchanged, Professor Sylvester gave the name of *Reciprocants*.

According to the general theory with respect to the inversion of the independent and dependent variable, we must have the relations:

$$\begin{aligned} \frac{dy}{dx} &= 1 \Big/ \frac{dx}{dy}, \frac{d^2y}{dx^2} = \frac{d}{dx} \left(1 \Big/ \frac{dx}{dy} \right) = \frac{d}{dy} \left(1 \Big/ \frac{dx}{dy} \right) \frac{dy}{dx} = - \frac{d^2x}{dy^2} \Big/ \left(\frac{dx}{dy} \right)^3; \\ \frac{d^3y}{dx^3} &= \frac{d}{dy} \left[- \frac{d^2x}{dy^2} \Big/ \left(\frac{dx}{dy} \right)^3 \right] \frac{dy}{dx} = - \left[\frac{dx}{dy} \cdot \frac{d^3x}{dy^3} - 3 \left(\frac{d^2x}{dy^2} \right)^2 \right] \Big/ \left(\frac{dx}{dy} \right)^5; \\ \frac{d^4y}{dx^4} &= - \left[\left(\frac{dx}{dy} \right)^2 \left(\frac{d^4x}{dy^4} \right) - 10 \frac{dx}{dy} \cdot \frac{d^2x}{dy^2} \cdot \frac{d^3x}{dy^3} + 15 \left(\frac{d^2x}{dy^2} \right)^3 \right] \Big/ \left(\frac{dx}{dy} \right)^7; \text{ etc.} \end{aligned}$$

After these relations are substituted for the various differential coefficients of y with respect to x , in any function of these differential coefficients or derivatives, we are said to have interchanged the independent and dependent variable.

Assume $dy \propto dx = T$, $d^2y \propto dx^2 = A \mid 2$, $d^3y \propto dx^3 = B \mid 3$, $d^4y \propto dx^4 = C \mid 4$, etc.; then, after eliminating the constants in the general equation of the straight line, by the method of differentiation, we obtain $d^2y \propto dx^2 = A$, $d^2x \propto dy^2 = 0$ (1).

The left-hand member of (1) is Professor Sylvester's first *pure* reciprocant, since it does not involve $dy \propto dx$; and this reciprocant is briefly and typically expressed by A . The third member of (1) represents the reciprocant when the independent and dependent variables x and y are interchanged.

The equation of the parabola $(\alpha x + \beta y)^2 + 2gx + 2fy + c = 0$, in which $\alpha^2 = a$ and $\beta^2 = b$, gives the second pure reciprocal:

$$3 \frac{d^2 y}{dx^2} \cdot \frac{d^4 y}{dx^4} - 5 \left(\frac{d^3 y}{dx^3} \right)^2 = 4AC - 5B^2, = 3 \frac{d^2 x}{dy^2} \cdot \frac{d^4 x}{dy^4} - 5 \left(\frac{d^3 x}{dy^3} \right)^2 = 0 \dots (2).$$

The general equation of a conic, in Cartesian co-ordinates, leads to the pure reciprocal:

$$\begin{aligned} & 9 \left(\frac{d^2 y}{dx^2} \right)^2 \cdot \frac{d^6 y}{dx^6} - 45 \frac{d^2 y}{dx^2} \cdot \frac{d^3 y}{dx^3} \cdot \frac{d^4 y}{dx^4} + 40 \left(\frac{d^3 y}{dx^3} \right)^3 = A^2 D - 3AB C + 2B^3, \\ & = 9 \left(\frac{d^2 x}{dy^2} \right)^2 \cdot \frac{d^6 x}{dy^6} - 45 \frac{d^2 x}{dy^2} \cdot \frac{d^3 x}{dy^3} \cdot \frac{d^4 x}{dy^4} + 40 \left(\frac{d^3 x}{dy^3} \right)^3 = 0 \dots (3), \text{ which is appropriately denominated the } \textit{Mongian Reciprocal}. \end{aligned}$$

Assume $xy = c$; then $xdy / dx + y = 0$, and

$$\begin{aligned} x = -2 \frac{dy}{dx} \int \frac{d^2 y}{dx^2} \dots \dots 2 \frac{dy}{dx} \cdot \frac{d^3 y}{dx^3} - 3 \left(\frac{d^2 y}{dx^2} \right)^2 = BT - A^2, = 2 \frac{dx}{dy} \cdot \frac{d^3 x}{dy^3} \\ - 3 \left(\frac{d^2 x}{dy^2} \right)^2 = 0 \dots (4), \text{ which is probably the simplest type of } \textit{mixed reciprocal}. \end{aligned}$$

From the equation of the hyperbola, $xy + ax + by + c = 0$, we can also deduce the *Schwarzian Reciprocal* represented by (4); and after dividing (4) by $(dy / dx)^2$, we have Professor Cayley's *Schwarzian Derivative*,

$$\frac{d^3 y / dx^3}{dy / dx} - \frac{3}{2} \left(\frac{d^2 y / dx^2}{dy / dx} \right)^2 = 0, \text{ which is directly deducible from } y = (ax + b)$$

$(Ax + B)$, and is of practical use in the solution of differential equations.

From the equation of the circle, $x^2 + y^2 = r^2$, we have $y(dy / dx) + x = 0$:

$$\begin{aligned} \left(\frac{dy}{dx} \right)^2 + y \frac{d^2 y}{dx^2} + 1 = 0; & 3 \frac{dy}{dx} \cdot \frac{d^2 y}{dx^2} \int \frac{d^3 y}{dx^3} + y = 0; 3 \left(\frac{d^2 y}{dx^2} \right)^2 \frac{d^3 y}{dx^3} \\ & - 3 \frac{dy}{dx} \cdot \frac{d^2 y}{dx^2} \cdot \frac{d^4 y}{dx^4} + 4 \left(\frac{d^3 y}{dx^3} \right)^2 \frac{dy}{dx} = A^2 B - 2(A C - B^2) T, \\ & = 3 \left(\frac{d^2 x}{dy^2} \right)^2 \frac{d^3 x}{dy^3} - 3 \frac{dx}{dy} \cdot \frac{d^2 x}{dy^2} \cdot \frac{d^4 x}{dy^4} + 4 \left(\frac{d^3 x}{dy^3} \right)^2 \frac{dx}{dy} = 0 \dots (5). \end{aligned}$$

The general equation of the circle, after three differentiations, gives

$$\begin{aligned} \left[1 + \left(\frac{dy}{dx} \right)^2 \right] \frac{d^3 y}{dx^3} - 3 \left(\frac{d^2 y}{dx^2} \right)^2 \frac{dy}{dx} &= (1 + T^2) B - 2A^2 T, \\ &= \left[1 + \left(\frac{dx}{dy} \right)^2 \right] \frac{d^3 x}{dy^3} - 3 \left(\frac{d^2 x}{dy^2} \right)^2 \frac{dx}{dy} = 0 \dots (6). \end{aligned}$$

The reciprocant represented by (6) may be written

$$1 + \left(\frac{dy}{dx}\right)^2 - 3 \left(\frac{d^2y}{dx^2}\right)^2 \frac{dy}{dx} \bigg/ \frac{d^3y}{dx^3} = 0 \dots (\alpha).$$

After differentiating (α) , etc., we have the reciprocant represented by (5).

From the equation $x^2 + xy + y^2 = 1$ may be deduced by differentiation, etc.,

$$x, = - \left(\frac{2dy + dx}{dy + 2dx} \right) y, = - \left(\frac{2(dy/dx + 1)}{dy/dx + 2} \right) y \dots (a);$$

$$\left(\frac{dy}{dx} + 2 \right)^2 = - \left(2 \frac{dy}{dx} + 1 \right) \left(\frac{dy}{dx} + 2 \right) \frac{dy}{dx} - 3y \frac{d^2y}{dx^2} \dots (b);$$

$$3y = - \left[2 \left(\frac{dy}{dx} \right)^3 + 6 \left(\frac{dy}{dx} \right)^2 + 6 \left(\frac{dy}{dx} \right) + 1 \right] \bigg/ \frac{d^2y}{dx^2} \dots (c).$$

$$\therefore 6 \left(\frac{dy}{dx} \right)^2 \left(\frac{d^2y}{dx^2} \right)^2 + 15 \left(\frac{d^2y}{dx^2} \right)^2 \frac{dy}{dx} + 6 \left(\frac{d^2y}{dx^2} \right)^2 - 2 \left(\frac{dy}{dx} \right)^3 \frac{d^3y}{dx^3} - 6 \left(\frac{dy}{dx} \right)^2 \frac{d^3y}{dx^3} \\ - 6 \frac{dy}{dx} \frac{d^3y}{dx^3} - 4 \frac{d^3y}{dx^3}, = T^3 - \left(\frac{2A^2 - 3B}{B} \right) T^2 - \left(\frac{5A^2 - 3B}{B} \right) T - 2 \left(\frac{A^2 - B}{B} \right), = 0$$

$\dots (7)$, which is a *cubic* reciprocant. From (6) is deduced $T^2 - 2(A^2/B)T + 1 = 0$, which is a *quadratic* reciprocant. Transforming (5), we have $T = A^2 B / 2(A^2 - B^2); = A^2/B, = 5AB/2C, = \text{etc.}$, which are *linear* reciprocants. In so far as their number is concerned, the pure reciprocants are like the major planets of the solar system—few; while the mixed reciprocants are like the minor planets of the same system—many.

NOTE.—Since we had the good fortune to be one of Professor Sylvester's students—one of the last and probably the youngest he had—in the Johns Hopkins University, we declare that Dr. Halsted's biographical sketch of that *enthusiastic* mathematical professor and investigator is the acme of appropriateness; and we also declare that all the commendation which that sketch accords to Professor Sylvester, is fully merited by him.

IS THE SUPPLEMENTARY ANGLE FINITE OR NOT?

By Professor JOHN N. LYLE, Ph. D., Westminster College, Fulton, Missouri.

If any individual angle whatever is greater than one right angle and